

A LIMIT SHAPE THEOREM FOR PERIODIC STOCHASTIC DISPERSION.

D. DOLGOPYAT, V. KALOSHIN AND L. KORALOV

ABSTRACT. We consider the evolution of a connected set on the plane carried by a periodic incompressible stochastic flow. While for almost every realization of the random flow at time t most of the particles are at a distance of order $\sqrt{t}$ away from the origin, there is a measure zero set of points, which escape to infinity at the linear rate. We study the set of points visited by the original set by time t , and show that such a set, when scaled down by the factor of t , has a limiting non random shape.

1. INTRODUCTION.

This paper deals with the long time behavior of a passive scalar carried by an incompressible random flow. As has been demonstrated for a large class of stochastic flows with zero mean, under some mixing conditions on the flow, the displacement of a single particle is typically of order $\sqrt{t}$ for large t . In [6] the authors show that for almost every realization of the random flow, if one considers the image of an open set under the action of the flow, then its spatial distribution, scaled by the square root of time, converges weakly to a Gaussian distribution. On the other hand, it has been shown in the work of Cranston, Scheutzow, Steinsaltz, and Lisei ([4], [5], [8], [10]) that in any open set there are points which escape to infinity at the linear rate. Denote the original set by Ω . One can think of Ω as of an oil spill or of a pollutant, say on the surface of the ocean. The evolution of the set under the action of the flow will be denoted by Ω_t .

We shall study the set of "poisoned" points, that is those visited by the image of Ω before time t

$$\mathcal{W}_t(\Omega) = \bigcup_{s \leq t} \Omega_s .$$

As shown in [4] and [5] the diameter of this set grows linearly in time almost surely. We shall be interested in its limit shape.

D.D. was partially supported by NSF and Sloan Foundation, V. K. was partially supported by American Institute of Mathematics Fellowship and Courant Institute, and L. K. was partially supported by NSF postdoctoral fellowship.

Consider a stochastic flow of diffeomorphisms on $\mathbb{R}^2$ generated by a finite-dimensional Brownian motion

$$(1) \quad dx_t = \sum_{k=1}^d X_k(x_t) \circ d\theta_k(t) + X_0(x_t)dt$$

where $X_0, X_1, \dots, X_d$ are C^∞ divergence free periodic vector fields and $\vec{\theta}(t) = (\theta_1(t), \dots, \theta_d(t))$ is a standard $\mathbb{R}^d$ -valued Brownian motion.

We impose several assumptions on the vector fields $X_0, X_1, \dots, X_d$, which are stated in the next section (cf [6]). All those, except the assumption of zero drift, are *nondegeneracy assumptions* and are satisfied for a generic set of vector fields $X_0, X_1, \dots, X_d$.

The main result of this paper is the following:

Theorem 1. (The Shape Theorem) *Let the original set Ω be bounded and contain a continuous curve. Under Assumptions (A)-(E) from Section 2 on the vector fields, there is a compact convex non random set $\mathcal{B}$, independent of Ω , such that for any $\varepsilon > 0$ almost surely*

$$(2) \quad (1 - \varepsilon)t\mathcal{B} \subset \mathcal{W}_t(\Omega) \subset (1 + \varepsilon)t\mathcal{B}$$

for all sufficiently large t .

In Section 3 we employ the estimates on the behavior of the characteristic function of a measure carried by the flow in order to bound from above the time it takes for the image of a curve to reach a fixed neighborhood of a far away point (cf [6]). This bound in turn implies the lower bound in (2) for the set $\mathcal{W}_t$.

The key element in the proof of the upper bound for $\mathcal{W}_t$ is to show that the set $\mathcal{W}_t$ for large t is almost independent of the original set (which, as will be demonstrated, can be taken to be a curve). In order to prove this we show that given two bounded curves γ and γ' we can almost surely find a contour, which contains γ' inside, and which consists of a finite number of pieces of γ_t , and a finite number of stable manifolds (whose length tends to zero as they evolve with the flow). In this way we see that if a point is visited by the image of γ' , then its small neighborhood is earlier visited by the image of γ .

In Section 4 we describe the construction of the contour, and in Section 5 we complete the proof of the upper bound.

2. NONDEGENERACY ASSUMPTIONS.

In this section we formulate a set of assumptions on the vector fields, which in particular imply the Central Limit Theorem for measures and the estimates on the behavior of the characteristic function of a measure carried by the flow (see [6]). Such estimates are essential in the proof

of the Shape Theorem. Recall that $X_0, X_1, \dots, X_d$ are assumed to be periodic and divergence free.

(A) (*hypoellipticity for x_t*) For all $x \in \mathbb{R}^2$ we have

$$\text{Lie}(X_1, \dots, X_d)(x) = \mathbb{R}^2.$$

Denote the diagonal in $\mathbb{T}^2 \times \mathbb{T}^2$ by

$$\Delta = \{(x^1, x^2) \in \mathbb{R}^2 \times \mathbb{R}^2 : x^1 = x^2 \pmod{1}\}.$$

(B) (*hypoellipticity for the two-point motion*) The generator of the two-point motion $\{(x_t^1, x_t^2) : t > 0\}$ is nondegenerate away from the diagonal Δ , meaning that the Lie brackets made out of $(X_1(x^1), X_1(x^2)), \dots, (X_d(x^1), X_d(x^2))$ generate $\mathbb{R}^2 \times \mathbb{R}^2$.

To formulate the next assumption we need additional notations. Let $Dx_t : T_{x_0}\mathbb{R}^2 \rightarrow T_{x_t}\mathbb{R}^2$ be the linearization of x_t at t . We need the hypoellipticity of the process $\{(x_t, Dx_t) : t > 0\}$. Denote by TX_k the derivative of the vector field X_k thought as the map on $T\mathbb{R}^2$ and by $S\mathbb{R}^2 = \{v \in T\mathbb{R}^2 : |v| = 1\}$ the unit tangent bundle on $\mathbb{R}^2$. If we denote by $\tilde{X}_k(v)$ the projection of $TX_k(v)$ onto $T_v S\mathbb{R}^2$, then the stochastic flow (1) on $\mathbb{R}^2$ induces a stochastic flow on the unit tangent bundle $S\mathbb{R}^2$ defined by the following equation:

$$d\tilde{x}_t = \sum_{k=1}^d \tilde{X}_k(\tilde{x}_t) \circ d\theta_k(t) + \tilde{X}_0(\tilde{x}_t)dt.$$

With these notations we have condition

(C) (*hypoellipticity for (x_t, Dx_t)*) For all $v \in S\mathbb{R}^2$ we have

$$\text{Lie}(\tilde{X}_1, \dots, \tilde{X}_d)(v) = T_v S\mathbb{R}^2.$$

For measure-preserving stochastic flows with conditions (C) Lyapunov exponents λ_1, λ_2 exist by *multiplicative ergodic theorem for stochastic flows* of diffeomorphisms (see [3], thm. 2.1). Moreover, the sum of Lyapunov exponents $\lambda_1 + \lambda_2$ should be zero (see e.g. [2]). Under conditions (A)–(C) the leading Lyapunov exponent is positive

$$\lambda_1 = \lim_{t \rightarrow \infty} \frac{1}{t} \log |Dx_t(x)(v)| > 0.$$

Indeed, Theorem 6.8 of [1] states that under condition (A) the maximal Lyapunov exponent λ_1 can be zero only if for almost every realization of the flow (1) one of the following two conditions is satisfied

- (a) there is a Riemannian metric $\mathbf{d}$ invariant with respect to the flow (1) or
- (b) there is a direction field $v(x)$ on $\mathbb{T}^2$ invariant with respect to the flow (1).

However (a) contradicts condition (B). Indeed, (a) implies that all the Lie brackets of $\{(X_k(x^1), X_k(x^2))\}_k$ are tangent to the leaves of the foliation

$$\{(x^1, x^2) \in \mathbb{T}^2 \times \mathbb{T}^2 : \mathbf{d}(x^1, x^2) = \text{Const} \}$$

and don't form the whole tangent space. On the other hand (b) contradicts condition (C), since (b) implies that all the Lie brackets are tangent to the graph of v . This positivity of λ_1 is crucial for our approach.

Let $L_{X_k}X_k(x)$ denote the derivative of X_k along X_k at the point x . Notice that $\sum_{k=1}^d L_{X_k}X_k + X_0$ is the deterministic components of the stochastic flow (1) rewritten in Ito's form. We require that the flow has no deterministic drift, which is expressed by the following conditions.

(D) (*zero drift*)

$$(3) \quad \int_{\mathbb{T}^2} \left(\sum_{k=1}^d L_{X_k}X_k + X_0 \right) (x) dx = 0 ,$$

$$(4) \quad \int_{\mathbb{T}^2} X_k(x) dx = 0 , \quad k = 1, \dots, d .$$

Assumption (4) is not needed for the Shape Theorem to be true. Still, in order to simplify the proof, that is in order to use the results of [6] without technical modifications, we shall assume (4) to hold.

The last assumption is concerned with the geometry of the stream lines of the vector fields $X_1, \dots, X_d$. Fix a coordinate system on the 2-torus $\mathbb{T}^2 = \{x = (x_1, x_2) \bmod 1\}$. As the vector fields have zero mean and are divergence free, there are periodic *stream* functions $H_1, \dots, H_d$, such that $X_k(x) = (-H'_{x_2}, H'_{x_1})$. Also, by condition (A), none of the points $x \in \mathbb{T}^2$ is a critical point for all of the stream functions $H_1, \dots, H_d$ at the same time. We require the following

(E) (*condition on the critical points of H_k*) For each k all of the critical points of H_k are non degenerate, and all the critical values are distinct.

3. LOWER BOUND.

3.1. An auxiliary statement. The following estimate will be repeatedly used in the proof of the lower bound.

Lemma 2. *Let $\{\xi_j\}$ be a sequence of random variables such that*

$$\mathbb{E}(\xi_{j+1} | \xi_1 \dots \xi_j) \leq 0,$$

and that for any N the sequence $\{\mathbb{E}|\xi_j|^N\}_j$ is bounded. Then for any $\varepsilon > 0$ there exists $\kappa = \kappa(\varepsilon, N) > 0$ such that for each $n \in \mathbb{Z}_+$ we have

$$\mathbb{P} \left\{ \sum_{j=1}^n \xi_j \geq \varepsilon n \right\} \leq \kappa n^{-N}.$$

Proof. Define a sequence of random variables $\zeta_j = \xi_j - \mathbb{E}(\xi_j | \xi_1 \dots \xi_{j-1})$. Then $M_n = \sum_{j=1}^n \zeta_j$ is a martingale whose quadratic variation is equal to $\langle M \rangle_n = \sum_{j=1}^n \zeta_j^2$. By the martingale inequality

$$\mathbb{E} M_n^{2N} \leq C_N \mathbb{E} \langle M \rangle_n^N \leq C'_N n^N.$$

Therefore, by the Chebyshev inequality

$$\mathbb{P} \left\{ \sum_{j=1}^n \zeta_j \geq \varepsilon n \right\} \leq \mathbb{P} \{ M_n^{2N} \geq (\varepsilon n)^{2N} \} \leq \kappa n^{-N}.$$

Since $\xi_j \leq \zeta_j$ the result also holds for the original sequence $\{\xi_j\}$. $\square$

3.2. Linear growth, an estimate from below. Let the initial set be a curve γ_0 , and consider a point A in the plane. We shall estimate the tail of the probability distribution of the time it takes for the curve to reach an R -neighborhood of A in terms of the distance between γ_0 and A (the constant R will be selected later). See Figure 1.

We shall call a curve *long* if its diameter is greater or equal than 1. Without loss of generality we may assume that the original curve is long. Given a curve γ_0 and a point A we define $\tau^R = \tau^R(\gamma_0, A)$ to be the first moment of time when the image of γ_0 reaches the R -neighborhood of A , and at the same time the image of γ_0 is long, that is

$$(5) \quad \tau^R = \inf \{ t > 0 : \text{dist}(\gamma_t, A) \leq R, \text{diam}(\gamma_t) \geq 1 \}.$$

Let $\mathcal{W}_t^R(\gamma_0)$ be the R -neighborhood of $\mathcal{W}_t(\gamma_0)$, that is the set of points, whose R -neighborhood is visited by the image of the original set before time t ,

$$\mathcal{W}_t^R(\gamma_0) = \{ x : \text{dist}(x, \gamma_s) \leq R \text{ for some } s \leq t \}.$$

Proposition 3. *Let $d = \text{dist}(A, \gamma_0)$. There is a constant $R > 0$, such that for any m there exist positive C and β such that for any long curve $\gamma_0 \subset \mathbb{R}^2$ and any point $A \in \mathbb{R}^2$ we have*

$$\mathbb{P} \{ \tau^R > \beta d \} \leq C d^{-m}.$$

The proof of Proposition 3 will rely on two other lemmas. We first state the lemma (proved in [5]), which shows that γ_t can not grow faster than linearly.

Lemma 4. ([5]) *Let γ_0 be the initial curve and*

$$\Phi_t = \sup_{x_0 \in \gamma_0} \sup_{0 \leq s \leq t} \|x_s - x_0\|.$$

(a) *There is a constant C such that almost surely*

$$\limsup_{t \rightarrow \infty} \frac{\Phi_t}{t} \leq C,$$

(b) *For any positive r and α we have*

$$\sup_{t \geq 1} \mathbb{E} \left(\exp \left[\frac{r \Phi_t^2}{t^2 \max(1, \ln(\Phi_t/t))^{2+\alpha}} \right] \right) < \infty.$$




FIGURE 1.

Given a pair (A_0, γ_0) , where γ_0 is a long curve, and A_0 is a point on γ_0 we define inductively the sequence (A_j, γ_j) as follows. Suppose that (A_j, γ_j) is defined so that

- γ_j is a connected interval of the image of γ_0 , i.e. $\gamma_j \subset \gamma_{t_j}$;
- $A_j \in \gamma_j$;
- γ_j is long.

Given $\alpha \in (0, \frac{\pi}{2})$ define the truncated α -cone

$$K_j(\alpha) = \{x : \text{dist}(x, A_j) \geq 1 \quad \text{and} \quad \angle(xA_jA) \leq \alpha\},$$

where $\angle(xA_jA)$ is the angle between the segments $[A_j, x]$ and $[A_j, A]$.

Let $\gamma_j(t)$ be the image of γ_j under the flow (so that $\gamma_j = \gamma_j(t_j)$.) Let t_{j+1} be the first moment such that

- $t_{j+1} - t_j \geq 1$;
- $\text{diam}(\gamma_j(t_{j+1})) \geq 1$;
- $\gamma_j(t_{j+1}) \cap K_j(\alpha) \neq \emptyset$.

Let A_{j+1} be an arbitrary point in $\gamma_j(t_{j+1}) \cap K_j(\alpha)$, $B_R(A_{j+1})$ denote the R -ball around A_{j+1} , and γ_{j+1} be a connected interval of $\gamma_j(t_{j+1}) \cap B_1(A_{j+1})$ containing A_{j+1} . Let $\mathcal{F}_{t_j}$ be the σ algebra of events determined prior to time t_j .

Lemma 5. *There exists $\alpha < \frac{\pi}{2}$ such that for any m we have*

$$\mathbb{E}((t_{j+1} - t_j)^m | \mathcal{F}_{t_j}) < C_m.$$

Proof. It is sufficient to prove that $\mathbb{E}(t_1^m) < C_m$ with C_m independent of the original long curve γ_0 . Without loss of generality we may assume that the point A_0 on the curve γ_0 coincides with the origin, and that the original curve is contained in a ball of radius 2 centered around the origin.

Fix some $0 < p < 1$. Since γ_0 is long, the Frostman Lemma ([9]) implies that there exists a probability measure ν on γ_0 whose p energy is bounded

$$(6) \quad I_p(\nu) = \int \int_{\gamma_0 \times \gamma_0} \frac{d\nu(x)d\nu(y)}{|x - y|^p} \leq C_p.$$

Let $f(x)$ be a continuous non-negative function compactly supported inside $K_0(\alpha)$. By the results of [6] under condition (6) there exists a non degenerate 2×2 matrix D such that for any $\rho, m > 0$ there exists T such that for all $t > T$

$$(7) \quad \mathbb{P} \left\{ \left| \int_{\gamma_0} f \left(\frac{x_t}{\sqrt{t}} \right) d\nu - \bar{f} \right| > \rho \right\} \leq t^{-m},$$

where $\bar{f}$ denotes the integral of f with respect to the Gaussian measure ν with zero mean and variance D . The last inequality of Sec. 7 in [6] establishes (7) for functions of the form $f(x) = \exp(i\xi x)$, but this inequality is also valid for compactly supported functions, since they could be uniformly approximated linear combinations of complex exponentials. Thus, for large t

$$\mathbb{P} \left\{ \gamma_t \cap \sqrt{t} \text{supp}(f) = \emptyset \right\} \leq t^{-m}.$$

Since for $t \geq 1$, we have that $\sqrt{t} \operatorname{supp}(f) \subset K_0(\alpha)$ we get that for $t \geq T$

$$\mathbb{P}\{\gamma_t \cap K_0(\alpha) = \emptyset\} \leq t^{-m}.$$

From (7) it also follows that for large t

$$\mathbb{P}\{\operatorname{diam}(\gamma_t) < 1\} \leq t^{-m}.$$

Since m is arbitrary, this implies the required result. $\square$

Proof of Proposition 3. Let $r_j = \operatorname{dist}(A_j, A)$. Let us show that there exists $R > 0$ such that for any m there is β for which

$$(8) \quad \mathbb{P}\{r_j > R \text{ for } j = 1, 2, \dots, [\beta r_0]\} \leq \frac{C_m}{r_0^m}.$$

We first establish a weaker bound: there is $R > 0$ such that

$$(9) \quad \mathbb{P}\{r_j > R \text{ for } j = 1, 2, \dots, [r_0^m]\} \leq \frac{C_m}{r_0^{m-1}}.$$

Note that there exist positive constants R and K such that if $r_0 > R$, then $\mathbb{E}(r_1 - r_0) \leq -K$. Let j_1 be the first moment when $r_j \leq R$, and in general let $\{j_n\}_n$ be the times of the consecutive visits of A_j to $B_R(A)$, that is $A_{j_n} \in B_R(A)$.

Then

$$(10) \quad r_{\min(j, j_1)} + K \min(j, j_1)$$

is a supermartingale. Hence

$$r_0 \geq \mathbb{E}(r_{\min([r_0^m], j_1)} + K \min([r_0^m], j_1)) \geq K[r_0^m] \mathbb{P}\{j_1 > [r_0^m]\},$$

which implies (9).

Recall the definition of the points A_j . Notice that for some constants C_1, C_2 and R_1 , which depend on α , we have if $\operatorname{dist}(A_{j+1}, A_j) \leq C_2 r_j$, then $r_{j+1} < r_j - R_1$. Indeed, A_{j+1} belongs to the truncated cone $K_j(\alpha)$ with the origin at A_j and directed toward A . This implies that the sequence $r_0, r_1, \dots, r_{[C_1 r_0]}$ contains an element r_j such that either $r_j \leq r_0^{\frac{1}{m}}$ or $\operatorname{dist}(A_{j+1}, A_j) > C_2 r_0^{\frac{1}{m}}$.

Let now j^* be the first moment when $r_j \leq r_0^{\frac{1}{m}}$. We have

$$\mathbb{P}\{j_1 \geq 2C_1 r_0\} \leq \mathbb{P}\{j^* > C_1 r_0\} + \mathbb{P}\{j_1 \geq 2C_1 r_0 | j^* \leq C_1 r_0\} = P_1 + P_2.$$

In order to estimate the first term on the right hand side of this inequality we note that if $j^* > C_1 r_0$ then $\operatorname{dist}(A_{j+1}, A_j) > C_2 r_0^{\frac{1}{m}}$ for some $j \leq [C_1 r_0]$. From Lemmas 4 and 5 it now follows that P_1 decays faster than any power of r_0 , in particular $P_1 \leq \frac{C_3}{\sqrt{r_0}}$.

We estimate the second term on the right hand side using (9) with $m = 2$

$$P_2 \leq C_3 r_0^{-\frac{m-1}{m}} = \frac{C_3}{\sqrt{r_0}}.$$

Therefore,

$$(11) \quad \mathbb{P}\{j_1 \geq 2C_1 r_0\} \leq \frac{C_4}{\sqrt{r_0}}.$$

Let a positive integer N be fixed. Then

$$\begin{aligned} \mathbb{P}\{j_1 \geq 2C_1 N r_0\} &= \mathbb{P}\{j_1 \geq 2C_1 N r_0; r_{[2C_1 r_0]} > r_0\} + \\ &\mathbb{P}\{j_1 \geq 2C_1 r_0; r_{[2C_1 r_0]} \leq r_0\} \mathbb{P}\{j_1 \geq 2C_1 N r_0 \mid j_1 \geq 2C_1 r_0; r_{[2C_1 r_0]} \leq r_0\} \\ &= Q_1 + Q_2 Q_3. \end{aligned}$$

By Lemmas 4 and 5 the tail of the probability distribution of $r_{j+1} - r_j$ decays faster than any power. From this fact and the fact that (10) is a supermartingale by Lemma 2 it is seen that Q_1 decays faster than any power of r_0 .

The factor Q_2 on the right hand side of the last inequality is estimated from above by $\frac{C_4}{\sqrt{r_0}}$ due to (11). Using the Markov property and continuing by induction we see that Q_3 is estimated from above by $C_5 r_0^{-\frac{N-1}{2}}$. We conclude that

$$\mathbb{P}\{j_1 \geq 2C_1 N r_0\} \leq C_6 r_0^{-\frac{N}{2}}.$$

Since N was arbitrary this implies (8).

Note that the estimate (8) remains valid even if the distance from the original point A_0 to A is less than r_0 . Therefore any $m > 0$ there are constants C_m and β such that

$$\mathbb{P}\{t_{j_1} > \beta d\} \leq \frac{C_m}{d^m}$$

for all d . By the Markov property

$$(12) \quad \mathbb{E}(t_{j_{n+1}} - t_{j_n} | \mathcal{F}_{t_{j_n}}) \leq C$$

for some C and for all n . We can therefore apply Lemma 2 with $\xi_n = t_{j_{n+1}} - t_{j_n} - C$ to obtain that for any N there exists κ such that

$$(13) \quad \mathbb{P}\{t_{j_n} > 2Cn\} \leq \kappa n^{-N}.$$

This combined with (8) completes the proof of the proposition. $\square$

Corollary 6. *There exist a positive constants c and R such that almost surely*

$B_{ct}(0) \subset \mathcal{W}_t^R$ for large t .

Proof. Consider a covering of $B_{ct}(0)$ by balls of radius $R/2$. By Proposition 3 for each of the balls of radius $R/2$, the probability that it is not visited by the curve by time t decays exponentially in t , provided that c is small enough, and R is large enough. On the other hand, for each c and R the number of balls needed to cover $B_{ct}(0)$ grows like t^2 times a constant. Therefore, the probability that the R neighborhood of some point in $B_{ct}(0)$ is not visited by the curve before time t decays exponentially in t . The Corollary follows by Borel-Cantelli Lemma. $\square$

From now on we fix R for which Proposition 3 and Corollary 6 hold.

Note that the bounds we obtained in the proof of Proposition 3 are uniform over all long curves. Let us employ this fact in the following corollary. Let $\mathcal{C}$ be the family of long curves which lie completely inside $B_{2R}(0)$ (we may assume that $R > 1$).

Corollary 7. *The family of stopping times*

$$\left\{ \frac{\tau^R(\gamma_0, tv)}{t} \right\}_{t \geq 1, \|v\|=1, \gamma_0 \in \mathcal{C}}$$

is uniformly integrable.

3.3. Stable norm. We shall now use the asymptotic of τ^R (the time it takes a curve to reach the R -neighborhood of a far away point) in order to define the limiting shape $\mathcal{B}$. Consider

$$|v|^R = \sup_{\mathcal{C}} \mathbb{E} \tau^R(\gamma, v).$$

Due to the stationarity of the underlying Brownian motion, and due to the periodicity of the vector fields we have

$$\mathbb{E} \tau^{2R}(\gamma, (t_1 + t_2)v) \leq \mathbb{E} \tau^R(\gamma, t_1 v) + \mathbb{E} \tau^R(\gamma_1, t_2 v),$$

where $\gamma_1 \in \mathcal{C}$ is some integer translation of a part of $\gamma_{\tau(\gamma, t_1 v)}$. Since by Proposition 3

$$\mathbb{E} \tau^R(\gamma, (t_1 + t_2)v) \leq \mathbb{E} \tau^{2R}(\gamma, (t_1 + t_2)v) + C$$

for some $C > 0$, it follows that the function $|tv|^R + C$ is sub-additive. Let $\|v\|^R = \lim_{t \rightarrow \infty} \frac{|tv|^R}{t}$. Similarly for $0 \leq s \leq 1$

$$|t(sv_1 + (1-s)v_2)|^R \leq |t sv_1|^R + |t(1-s)v_2|^R + C,$$

so

$$\|sv_1 + (1-s)v_2\|^R \leq s\|v_1\|^R + (1-s)\|v_2\|^R.$$

Let $\mathcal{B} = \{v : \|v\|^R \leq 1\}$. By the remarks above and by Corollary 7 we have that $\mathcal{B}$ is a convex compact set. It will be seen that the norm $\|v\|^R$ and the set $\mathcal{B}$ are independent of R .

Lemma 8. *For any $\varepsilon > 0$ almost surely $(1 - \varepsilon)t\mathcal{B} \subset \mathcal{W}_t^R$ for large enough t .*

Proof. It suffices to show that for any v and N

$$\mathbb{P}\{\tau^R(\gamma, tv) \geq (1 + \varepsilon)t\|v\|^R\} \leq C_N t^{-N}.$$

By the definition of $\|v\|^R$ there exists t_0 such that

$$(14) \quad \mathbb{E} \tau^R(\gamma, tv) \leq t(1 + \frac{\varepsilon}{3})\|v\|^R$$

for any $t \geq t_0$ and $\gamma \in \mathcal{C}$. Define the stopping time τ_1 as follows

$$\tau_1 = \inf\{t : \gamma_t \cap B_R(t_0 v) \neq \emptyset; \text{diam}(\gamma_t) \geq 1\}.$$

Let $\gamma^{(1)}$ be a part of γ_{τ_1} , which is long, is contained in $B_{2R}(t_0 v)$, and has a non empty intersection with $B_R(t_0 v)$. Similarly we define τ_2 to be the first time following τ_1 , when the image of $\gamma^{(1)}$ is long and intersects $B_R(2t_0 v)$, and let $\gamma^{(2)}$ be a part of the image of $\gamma^{(1)}$, and so on. We have therefore constructed a sequence of stopping times, such that

$$\tau^R(\gamma, nt_0 v) \leq \sum_{j=1}^n (\tau_j - \tau_{j-1}).$$

By (14), due to the periodicity of the underlying vector fields, for large enough t_0 we have

$$\mathbb{E} (\tau_j - \tau_{j-1}) \leq t_0 \left(1 + \frac{2\varepsilon}{3}\right) \|v\|^R.$$

Now the result follows by Lemma 2 and Proposition 3. $\square$

Now we prove that Lemma 8 remains valid even when the R -neighborhood of $\mathcal{W}_t$ is replaced by $\mathcal{W}_t(\gamma)$ itself.

Theorem 9. *For any $\varepsilon > 0$ we have almost surely $(1 - \varepsilon)t\mathcal{B} \subset \mathcal{W}_t(\gamma)$ for large enough t .*

This Theorem is a consequence of Proposition 3 and the fact that when a long curve reaches an R -neighborhood of a point, the distribution of the time it then takes for the curve to sweep the entire neighborhood has fast decreasing tail. Thus Theorem 9 follows from the standard Borel-Cantelli arguments and the following

Lemma 10. *Let γ_0 be a long curve such that $\text{dist}(\gamma_0, A) \leq R$. Let*

$$\sigma = \inf\{t > 0 : B_R(A) \subset \bigcup_{s \leq t} \gamma_s; \text{diam}(\gamma_t) \geq 1\}.$$

Then for any $m > 0$ and some C_m we have

$$(15) \quad \mathbb{P}(\sigma > t) \leq C_m t^{-m} .$$




FIGURE 2. The image of a sufficiently long curve γ covers a small ball $B_\delta(\tilde{A})$ with positive probability. (Dashed lines on the picture show the orbits of the endpoints of γ .)

Proof. Since the ball $B_R(A)$ can be covered by a finite number of arbitrarily small balls, it is sufficient to demonstrate (15) for

$$\sigma_1 = \inf \left\{ t : B_\delta(\tilde{A}) \subset \bigcup_{s \leq t} \gamma_s ; \text{diam}(\gamma_t) \geq 1 \right\} ,$$

where $\tilde{A} \in B_R(A)$. As before, let j_n be the times of the consecutive visits of A_j to $B_R(A)$. We claim that for small enough δ there exists $\theta < 1$ such that

$$(16) \quad \mathbb{P}\{\sigma_1 \geq t_{j_n} | \sigma_1 \geq t_{j_{n-1}}\} \leq \theta .$$

Since $t_{j_n} - t_{j_{n-1}} \geq 1$ it is sufficient to show that any point $\tilde{A}$ with probability $1 - \theta$ has a neighborhood, which is swept by the time 1 evolution of any long curve γ_0 , which initially is at a distance no greater than $2R$ away from $\tilde{A}$.

Due to condition (A) we may assume that $\tilde{A}$ is not a critical point of say X_1 . Let Γ be the streamline of X_1 , which passes through $\tilde{A}$. Then, due to the periodicity, Γ is either a closed curve, or an unbounded periodic curve, or $\tilde{A}$ belongs to a separatrix (level set containing a saddle point) of H_1 .

Consider the case when Γ is a closed curve. Let $U^{2\delta}$ be the region formed by the streamlines of X_1 passing through the 2δ neighborhood of $\tilde{A}$. For δ small enough the region $U^{2\delta}$ is homeomorphic to an annulus. Let B^1 and B^2 be the interior and the exterior circles bounding $U^{2\delta}$, and let P be a point inside B^1 . Define the winding number of a point x on a curve γ as the change of argument of rotation around P for the part of γ from the initial point to x . Due to the hypoellipticity of the two point motion with probability $p_1 > 0$ we have the following: one of the points on $\gamma_{\frac{1}{2}}$ lies inside B^1 , while another point on $\gamma_{\frac{1}{2}}$ lies outside B^2 , and the winding number around P is bounded in absolute value by a constant K for all points of $\gamma_{\frac{1}{2}}$. Notice, that in this case $\gamma_{\frac{1}{2}}$ crosses each of the streamlines, which form $U^{2\delta}$.

Let $\phi_t^1(x)$ be the evolution of x under the action of the vector field X_1 :

$$\dot{\phi}_t^1(x) = X_1(\phi_t^1(x)) , \quad \phi_0(x) = x .$$

Let T be the supremum over $U^{2\delta}$ of the time it takes for a point to make a complete rotation

$$T = \sup_{x \in U^{2\delta}} \inf_{t > 0} \{t : \phi_t^1(x) = x\} .$$

Given the event that $\gamma_{\frac{1}{2}}$ crosses each of the streamlines of $U^{2\delta}$, for any $\varepsilon > 0$ there is a positive probability $p_2 > 0$ that the time $t = \frac{1}{2}$ dynamics of (1) is ε -close to the solution of

$$(17) \quad \dot{\phi}_t^1(x) = 4T(K+1)X_1(\phi_t^1(x)) , \quad \phi_0^1(x) = x .$$

By selecting ε small enough we see that with positive probability p_2 the time $\frac{1}{2}$ evolution of $\gamma_{\frac{1}{2}}$ sweeps the entire annulus U^δ . Therefore, the probability that $B_\delta(\tilde{A})$ is swept by the image of γ_0 before $t = 1$ is not less than $p_1 p_2 > 0$. We can take $\theta = 1 - p_1 p_2$. The cases when Γ is a periodic unbounded curve, and when $\tilde{A}$ belongs to a separatrix of H_1 are treated similarly. (In the case when $\tilde{A}$ belongs to the separatrix we may need to use one of the vector fields $X_2, \dots, X_d$, or one of the commutators in order to show that the evolution of γ_0 may pass through a saddle point).

From (16) it follows by induction that

$$(18) \quad \mathbb{P}\{\sigma_1 \geq t_{j_n}\} \leq \theta^n$$

for some $\theta < 1$. This combined with estimate (13) completes the proof of the lemma. $\square$

3.4. Occupation times. The next result is used in Subsection 4.1.

Lemma 11. *For any $\varepsilon > 0$ there is a constant R^* such that for any curve γ for any A*

$$\liminf_{T \rightarrow \infty} \frac{|\{t \leq T : \gamma_t \cap B_{R^*}(A) \neq \emptyset\}|}{T} \geq 1 - \varepsilon.$$

Proof. Let j_n be as in Proposition 3. Let

$$a_n(R^*) = |\{t > 0 : t_{j_{n-1}} \leq t < t_{j_n} \quad \text{and} \quad A_{j_{n-1}}(t - t_{j_{n-1}}) \notin B_{R^*}(A)\}|$$

be the amount of time between two consecutive visits of a smaller R -ball $B_R(A)$ spend outside a bigger R^* -ball $B_{R^*}(A)$. Then finiteness of expectation of time (12) between consecutive visits of $B_R(A)$ implies that for each ε there is a sufficiently large R^* such that for all n

$$\mathbb{E} a_n(R^*) < \frac{\varepsilon}{2}.$$

Using Lemma 2 we obtain that almost surely $\sum_{n=1}^N a_n(R^*) < \varepsilon N$ for a large N . On the other hand by the definition of t_j

$$\sum_{n=1}^k (t_{j_n} - t_{j_{n-1}}) > k.$$

This implies the required result. $\square$

Corollary 12. *There is a constant R^* such that for any two open sets U_1, U_2 , any two points A_1, A_2 , and some T almost surely there is a moment $t > T$ such that $\phi_t U_j$ intersects $B_{R^*}(A_j)$ for both $j = 1, 2$.*

4. GEOMETRY OF STABLE MANIFOLDS.

4.1. Transitivity of stable lamination. Before we can proceed further we need to establish some properties of unstable lamination. Recall that one of our assumptions is $\lambda_1 > 0$. Let $0 < \bar{\lambda}_1 < \lambda_1$ then by stable manifold theorem [7] for all t almost surely for almost every x the set

$$W^s(x, t) = \{y : \text{for some } C(y) \text{ we have } d(f_{t,s}y, f_{t,s}x) \leq C(y)e^{-\bar{\lambda}_1(s-t)}\}$$

is a smooth curve passing through x . By stationarity the parameters of W^s such as its length, curvature etc. have distributions independent of t . Here we prove the following.

Theorem 13. *Almost surely $W^s(\cdot, t)$ is transitive for all t , that is, given two open sets U_1 and U_2 for some $x \in U_1$ we have $W^s(x, t) \cap U_2 \neq \emptyset$ or U_1 and U_2 are connected by a stable manifold.*

Proof. Since $W^s(\cdot, t) = \varphi_t W^s(\cdot, 0)$ and the image of a transitive lamination is transitive, it is enough to show that $W^s(\cdot, 0)$ is transitive. Let E_{U_1, U_2} denote the event

$$\{\{\theta(t)\}_{t>0} : \text{for some } y \ W^s(y, 0) \cap U_1 \neq \emptyset \text{ and } W^s(y, 0) \cap U_2 \neq \emptyset\}$$

We want to show that $\mathbb{P}\{E_{U_1, U_2}\} = 1$. In fact a weaker statement suffices.

Proposition 14. *Let $\mathcal{F}_t$ denote the σ -algebra generated by $\{\theta_j(s) | s \leq t\}$. Then for any $t > 0$ there is a constant $\delta > 0$ such that*

$$\mathbb{P}\{E_{U_1, U_2} | \mathcal{F}_t\} \geq \delta.$$

The rest of this subsection is devoted to the proof of Proposition 14. This proposition implies the Theorem because $\mathbb{P}\{E_{U_1, U_2} | \mathcal{F}_t\}$ converges as $t \rightarrow +\infty$ to the indicator of E_{U_1, U_2} almost surely and so we get that this indicator is greater than $\delta/2$ almost surely. Since the indicator assumes only values 0 and 1 Theorem 13 follows. $\square$

Lemma 15. *For any point $x \in \mathbb{R}^2$, positive r_0 , ε there exists $p_1 > 0$ such that for all $y \in \mathbb{R}^2$ such that $d(x, y) \leq r_0$ we have*

$$\mathbb{P}\{W^s(x, 0) \cap B(y, \varepsilon) \neq \emptyset\} > p_1.$$

Proof. There are positive r_1, p_1 such that for each pair: a point $x \in \mathbb{R}^2$ and a time $t > 0$ we have $\mathbb{P}\{\text{diam}(W^s(x, t)) > r_1\} > p_1$ or probability to have a stable manifold at a point at a certain time of diameter at least a positive r_1 . Thus for each positive ε_1 there is a constant $p_2 > 0$ such that for $x \in \mathbb{R}^2$ there is a point $y \in \mathbb{R}^2$ such that $d(x, y) = r_1/2$ and for all t we have $\mathbb{P}\{W^s(x, t) \cap B(y, \varepsilon_1) \neq \emptyset\} > p_2$. Another way to put it is that if a stable manifold at a point at time t is of diameter at least r_1 , then it intersects a small ball centered at a nearby point with a positive probability. Now by hypoellipticity of the two-point motion there are positive ε_1 and p_3 such that for each pair of points $\bar{x}, \bar{y} \in \mathbb{R}^2$ and $d(x, y) \leq r_0$ for the time 1 map of the flow (1) we have

$$\mathbb{P}\{\varphi_1 B(y, \varepsilon) \supset B(\bar{y}, \varepsilon_1) | x_1 = \bar{x}\} > p_3.$$

However, $W^s(x_1, 1)$ crosses $\varphi_1(B(y, \varepsilon))$ with positive probability, therefore,

$$\mathbb{P}\left\{W^s(x, t) \cap B(y, \varepsilon) \neq \emptyset\right\} > p_2 p_3 > 0.$$

This completes the proof. $\square$




FIGURE 3. A stable manifold joining $\mathcal{Q}_1$ and $\mathcal{Q}_2$ should intersect the image of U_1 unless it is long enough to avoid the tongue $\varphi_{t,t+\varepsilon}\Lambda$

Proof of Proposition 14. Fix $1 \leq k \leq d$ and consider the vector field X_k along with its stream function $H_k : \mathbb{T}^2 \rightarrow \mathbb{R}$. By condition (E) H_k exists and its lift $\hat{H}_k : \mathbb{R}^2 \rightarrow \mathbb{R}$ has unbounded level sets $\hat{H}_k^{-1}(c^*)$ for some regular values c^* . Choose two adjacent critical values c^-, c^+ of H_k with $c^- < c^* < c^+$. Then the strip of level sets $\mathcal{H}_k = \{\hat{H}_k^{-1}(c^-, c^+)\}$ consists of only unbounded level sets of $\hat{H}_k$ which are also stream lines of X_k . Let $\mathcal{Q}_1$ and $\mathcal{Q}_2$ be the components consisting of bounded streamlines bounding $\mathcal{H}_k$ (see figure 3). Choose $R^* > 0$ from Corollary 12. Combining Lemma 11 and Corollary 12 we have that given two point A_1, A_2 inside $\mathcal{Q}_1$ for each T there exists $t > T$ such that both sets $\varphi_t U_i \cap B_{R^*}(A_i) \cap \mathcal{Q}_j = \emptyset$ for all $i = 1, 2$ and $j = 1, 2$. In other words, the image of U_1 (resp. U_2) at time t bisects the strip $\mathcal{H}_k$ into two parts and we can restrict this *nonempty* intersection to not a long in the horizontal direction strip each. Moreover, we shall need both intersections to be well separated in the horizontal direction inside $\mathcal{H}_k$, or $\text{dist}(A_1, A_2) = l \gg R^*$.

Without the loss of generality one can assume that the points in $\mathcal{H}_k$ move in the direction from A_1 to A_2 under the vector field X_k . Take two small constants $\epsilon, \tau > 0$. Then with positive probability the dynamics of $\varphi_{t,t+\epsilon}$ is C^0 -close to the dynamics of $\Phi_{1/\tau}^k$, which is the time one

map of the flow generated by X_k/τ . Define $\Lambda^k = B_{R^*}(A_1) \cap \mathcal{H}_k$ and $\mathcal{T}_\tau^k = \Phi_{1/\tau}^k \Lambda^k$. Then with positive probability $\varphi_{t,t+\epsilon}(\mathcal{T}_\tau^k)$ is C^0 -close to $\mathcal{T}_\tau^k$. Moreover, $\varphi_{t+\epsilon}(U_1)$ divides it into two parts. On the other hand, since $\varphi_{1/\tau}$ does not move the set $\mathcal{Q}_1$, we know that the set $\varphi_{t+\epsilon}(U_2)$ still contains a point q in $\mathcal{Q}_1 \cap B_{R^*}(A_2)$. Now by Lemma 15 there is a positive probability that $W^s(q, t+\epsilon)$ intersects $\mathcal{Q}_2$, moreover, since R^* is sufficiently large, the segment of $W^s(q, t+\epsilon)$ between q and the first crossing of $\mathcal{Q}_2$ lies inside $B_{R^*}(A_2)$. But if l and $1/\tau$ are much larger than R^* this segment must cross the tongue $\Phi_{1/\tau}(\Lambda)$. $\square$

4.2. Fences. The following property of unstable foliations plays the key role in our analysis. Let $\mathcal{S}_N$ denote $N \times N$ square centered at 0.




FIGURE 4. The construction used in the proof of transitivity of the stable lamination also gives the existence of fences.

Theorem 16. *For any curve γ and any $N > 0$ there almost surely exist a time t and contour Γ consisting of two pieces of translated images of γ ($\gamma_t + \vec{n}_1$ and $\gamma_t + \vec{n}_2$), and two stable curves, $W^s(x_1, t)$ and $W^s(x_2, t)$, such that $\text{Int}(\Gamma)$ contains $\mathcal{S}_N$.*

Proof. Let X_k be the vector field with unbounded stream lines. Let $\mathcal{H}_k$ be a strip in $\mathbb{R}^2$, which contains some of the unbounded stream lines of X_k . We can then find two parallel translations of $\mathcal{H}_k$ by integer vectors, to be denoted $\mathcal{H}_k^1$ and $\mathcal{H}_k^2$, which contain the square $\mathcal{S}_N$ between them.

Thus, the plane is a union of five regions: the parallel strips $\mathcal{H}_k^1$ and $\mathcal{H}_k^2$, the strip between them, to be denoted by $\mathcal{R}$, which contains $\mathcal{S}_N$, and two half planes, $\mathcal{P}_1$, and $\mathcal{P}_2$.

Let $\varepsilon > 0$ be fixed. By Theorem 13, for any $t \geq 0$, there almost surely exists a stable manifold $W^s(x, t)$, which intersects both half planes, $\mathcal{P}_1$ and $\mathcal{P}_2$. By periodicity there are two stable manifolds, $W^s(x_1, t)$ and $W^s(x_2, t)$, such that $\mathcal{S}_N$ is contained in the part of $\mathcal{R}$ between $W^s(x_1, t)$ and $W^s(x_2, t)$.

Let r be such that $\text{diam}(W^s(x_1, t) \cup W^s(x_2, t) \cup \mathcal{S}_N) \leq r$ with probability at least $1 - \varepsilon$. Then by the hypoellipticity of the two point motion, and the recurrence condition (Proposition 3), with probability at least $1 - 2\varepsilon$ the curve γ contains a part $\hat{\gamma}$, whose image $\hat{\gamma}_t$ belongs to $\mathcal{H}_k^1$, and intersects any curve α connecting $\mathcal{P}_1$ with $\mathcal{P}_2$, with the property $\text{diam}(\alpha \cup \mathcal{S}_N) \leq r$. We also require the same of the integer translation of $\hat{\gamma}_t$ to H_2 .

The desired contour then consists of $\hat{\gamma}_t$, its integer translation to H_2 , and the stable curves $W^s(x_1, t)$ and $W^s(x_2, t)$. $\square$

We say that T is a *rearrangement* of $\mathcal{S}_N$ if there is a partition $T = \bigcup_j T_j$ of T into finitely many pieces and a bijection $\beta : T \rightarrow \mathcal{S}_N$ such that on each T_j we have $\beta(x) = x + \vec{n}_j$, where $\vec{n}_j \in \mathbb{Z}^2$.

From Theorem 16 it follows that there exists a contour Γ_0 , which consists of two pieces of translations of γ ($\gamma + \vec{n}_1$ and $\gamma + \vec{n}_2$), and two stable curves, $W^s(x_1, 0)$ and $W^s(x_2, 0)$, such that $\text{Int}(\Gamma_0)$ contains a rearrangement of $\mathcal{S}_N$.

5. UPPER BOUND.

5.1. Estimates in Probability. We first establish the asymptotic of the expectation of $\tau^R(\gamma, tv)$.

Lemma 17. *The following limit*

$$\lim_{t \rightarrow \infty} \frac{\mathbb{E} \tau^R(\gamma, tv)}{t} = \|v\|^R$$

is uniform in $\gamma \in \mathcal{C}$.

Proof. Let $S = [-3R, 3R]^2$. Then $\gamma \subset \text{Int}(S)$ and so $\tau^R(\gamma, tv) \geq \tau^R(S, tv)$. Hence it suffices to show that

$$\frac{\mathbb{E} \tau^R(S, tv)}{t} \rightarrow \|v\|^R.$$

By Corollary 7 it suffices to show that

$$\{\tau^R(\gamma, tv) - \tau^R(S, tv)\}_{\gamma \in \mathcal{C}, t \geq 1, \|v\|=1}$$

is tight. To prove the tightness let G be the set of all γ -fences containing a rearrangement of S . For $\Gamma \in G$ let $S'(\Gamma)$ be some rearrangement of S inside Γ and let $\sigma(\Gamma)$ be the part of Γ consisting of stable manifolds.

Then for any $\varepsilon > 0$ there exists r such that

(19)

$$\mathbb{P}\{\exists \Gamma \in G \text{ such that } \text{diam}(\Gamma \bigcup S) < r \text{ and } \sup_{t \geq 0} l(\sigma_t) < r\} \geq 1 - \varepsilon ,$$

Since by (19) the distance between any two points, first one from S , and the second one from S' , does not exceed r , and since S' is contained in Γ , we have that with probability not smaller than $1 - \varepsilon$ for any A there is the inequality $\tau^{r+R}(\Gamma, A) \leq \tau^R(S, A)$. Now (19) implies that $\text{dist}(\gamma_{\tau^{r+R}(\Gamma, A)}, A) \leq 2r + R$ since $B(A, r + R)$ intersects either $\gamma_{\tau^{r+R}(\Gamma, A)}$ or its integer translation by distance no larger than r , in which case $\text{dist}(\gamma_{\tau^{r+R}(\Gamma, A)}, A) \leq r + R$, or $B(A, r + R)$ intersects the stable curve, and since the length of this curve is at most r , there is a point in $\gamma_{\tau^{r+R}(\Gamma, A)}$ in $B(A, 2r + R)$. Now by Proposition 3 there is t such that

$$\mathbb{P}\{\tau^R(\gamma_{\tau^{r+R}(\Gamma, A)}, A) > t\} \leq 2\varepsilon.$$

This completes the proof. $\square$

Corollary 18. *For any $\gamma \in \mathcal{C}$ we have $\lim_{t \rightarrow \infty} \frac{\tau^R(\gamma, tv)}{t} = \|v\|^R$ in probability.*

Proof. By Corollary 7 the family $\{\frac{\tau^R(\gamma, tv)}{t}\}$ is tight. Let ν be some limit distribution of this family. By Lemma 8 $\text{supp } \nu \subset [\|v\|^R, \infty[$. By Lemma 17 $\int t d\nu(t) = \|v\|^R$. Thus $\nu = \delta_{\|v\|^R}$. $\square$

Corollary 19. *For any $\varepsilon > 0$*

$$\lim_{t \rightarrow \infty} \mathbb{P}\{\mathcal{W}_t^R \subset (1 + \varepsilon)t\mathcal{B}\} = 1 .$$

Proof. We need to show that

$$(20) \quad \mathbb{P}\{\exists v : \frac{\tau^R(\gamma, tv)}{t} < \frac{\|v\|^R}{1 + \varepsilon}\} \rightarrow 0.$$

Take $\varepsilon' \ll \varepsilon$. Let $\{v_j\}$ be an ε' -net in $\partial\mathcal{B}$. Then by Corollary 18

$$(21) \quad \mathbb{P}\left\{\exists j : \frac{\tau^R(\gamma, tv_j)}{t} < \frac{\|v_j\|^R}{1 + \frac{\varepsilon}{2}}\right\} \rightarrow 0.$$

However by Proposition 3 for any m for sufficiently small ε' we have

(22)

$$\mathbb{P}\{\exists j : \frac{\tau^R(\gamma, tv_j)}{t} < \frac{\|v_j\|^R}{1 + \frac{\varepsilon}{2}} | \exists v : \frac{\tau^R(\gamma, tv)}{t} < \frac{\|v\|^R}{1 + \varepsilon}\} \geq 1 - C_m t^{-m}.$$

Now, (21) and (22) combined yield (20). $\square$

5.2. Point-to-line passage. Given a line l let

$$\rho(l, t) = \sup_c \frac{\mathbb{E} (\tau^R(\gamma, tl))}{t}.$$

The following results are proved similarly to a point-to-point case.

Lemma 20. (a) $\frac{\mathbb{E} (\tau^R(\gamma, tl))}{t}$ converges uniformly to $\rho(l) = \lim_{t \rightarrow \infty} \rho(l, t)$.
 (b) For any $\varepsilon > 0$

$$\mathbb{P}\{\tau^R(\gamma, tl) > (\rho(l) + \varepsilon)t\} \leq C_N t^{-N}.$$

(c) $\frac{\tau^R(\gamma, tl)}{t} \rightarrow \rho(l)$ in probability.

The next lemma relates $\rho(l)$ to the norm $\|v\|^R$.




FIGURE 5. Line-to-line passage time differs little from square-to line-passage time since all squares reach the lines at about the same time

Lemma 21. For any line l

$$\rho(l) = \inf_{v \in l} \|v\|^R.$$

Proof. This follows from Lemma 20, Lemma 8 and Corollary 19. $\square$

Consider now

$$\rho^*(l) = \lim_{t \rightarrow \infty} \frac{\mathbb{E}(\tau^R(l', tl))}{t}$$

where l' is the line parallel to l passing through origin.

Lemma 22. *For any line l we have the equality*

$$\rho^*(l) = \rho(l).$$

Proof. Let Π be the union of fundamental domains containing l' . Applying the reasoning of subsection 5.1 we get

$$\mathbb{E} \left(\frac{\tau^R(l', tl) - \tau^R(\Pi, tl)}{t} \right) \rightarrow 0.$$

Let S be one fundamental domain from Π . Then

$$\mathbb{E} \left(\frac{\tau^R(S, tl) - \tau^R(\Pi, tl)}{t} \right) \rightarrow 0.$$

Subtracting we get

$$\mathbb{E} \left(\frac{\tau^R(l', tl) - \tau^R(S, tl)}{t} \right) \rightarrow 0.$$

Passing to the limit as $t \rightarrow \infty$ we get the statement required. $\square$

5.3. Almost sure convergence.

Theorem 23. *For any $\varepsilon > 0$ almost surely $\mathcal{W}_t^R \subset (1 + \varepsilon)t\mathcal{B}$ for large enough t .*

Proof. Again choose some $\varepsilon_1 \ll \varepsilon$ and let $\{v_j\}$ be an ε_1 -net on $\partial\mathcal{B}$. Let $\mathcal{B}_{\varepsilon_1}$ be the region bounded by support lines of $\mathcal{B}$ passing through $\{v_j\}$. It suffices to prove that almost surely $\mathcal{W}_t^R \subset t(1 + \varepsilon)\mathcal{B}_{\varepsilon_1}$ for large t . But this accounts to showing that almost surely

$$(23) \quad \tau^R(\gamma, tl) \geq (1 - \varepsilon)t\rho(l)$$

for large t . Let t^* be such that

$$(24) \quad \mathbb{E}(\tau^R(l', tl)) > (1 - \frac{\varepsilon}{2})(t + 1)\rho(l)$$

for $t > t^*$. Let $l_j = jt^*l$. Let Δ_j be the first moment after $\tau^R(\gamma, l_j)$ such that $l_j(\tau^R(\gamma, l_j) + \Delta) \cap l_{j+1} \neq \emptyset$. It follows that $\tau^R(\gamma, l_j) \geq \sum_{k=0}^{j-1} \Delta_k$. Thus (23) follows by (24) and Lemma 2. $\square$

REFERENCES

- [1] P. Baxendale *Lyapunov exponents and relative entropy for a stochastic flow of diffeomorphisms*, Probab. Th. & Rel. Fields **81** (1989) 521–554.
- [2] P. Baxendale, D. W. Stroock, *Large deviations and stochastic flows of diffeomorphisms*, Prob. Th. & Rel. Fields **80**, 169–215, (1988);
- [3] A. Carverhill, *Flows of stochastic dynamical systems: ergodic theory*, Stochastics, **14**, 273–317, (1985);
- [4] M. Cranston, M. Schuetzow & D. Steinsaltz, *Linear expansion of isotropic Brownian flows*, El. Comm. Prob., **4**, (1999); 91-101;
- [5] M. Cranston, M. Schuetzow & D. Steinsaltz. *Linear bounds for stochastic dispersion*, Ann. Prob., **28**, (2000), 1852–1869;
- [6] D. Dolgopyat, V. Kaloshin & L. Korolov, *Sample Path Properties of the Stochastic Flows*, preprint.
- [7] Y. Kifer, *Random perturbations of dynamical systems*, Progress in Probability and Statistics, 16. Birkhuser Boston, Inc., Boston, MA, 1988.
- [8] H. Lisei & M. Scheutzow *Linear bounds and Gaussian tails in a stochastic dispersion model*, preprint;
- [9] P. Mattila, *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability*. Cambridge Studies in Advanced Mathematics, 44. Cambridge University Press, Cambridge, 1995.
- [10] M. Schuetzow, & D. Steinsaltz, *Chasing balls through martingale fields*, preprint;